

Tooba Batool¹, Safa Toqir², Komal Naz³, Amina Sajjad⁴, Asma Saeed⁵, Mohammad Arsalan Aslam⁶, Farha Younas⁷, Misbah Nawaz⁸, Saba Mehnaz⁹ and Tabassam Fatima¹⁰

ABSTRACT

The recent growth of infectious diseases stemming from zoonotic origins has placed a significant global burden on both morbidity and mortality. Additionally, it has generated substantial economic challenges. The complexity and dynamism of the resurgence and epidemiology of zoonoses are shaped by diverse factors which can be broadly classified into human-related, pathogen-related and climate related parameters. The diagnosis of animal diseases serves a dual role as both the origin and solution to various ailments. It is instrumental in managing and preventing diseases, playing a crucial part in overall disease control. Therefore, the imperative for the development of veterinary diagnostic tools becomes apparent, ensuring comprehensive animal welfare and effective monitoring of disease spread. Here we discussed various molecular and non-molecular diagnostic methods for zoonotic infections. These diverse approaches include viral metagenomics, clinical recognition, standard laboratory assessments, and laboratory tests. By examining recent advancements in diagnostic methodologies, this chapter aims to underscore the importance of ongoing research and innovation in the field of zoonotic disease diagnostics for enhanced public health preparedness and intervention strategies.

Keywords: Disease management, Diagnostic, Zoonotic, Viral metagenomics, Clinical tests, PCR, DNA fingerprinting, DNA based composition, Radioimmunoassay.

CITATION

Batool T, Toqir S, Naz K, Sajjad A, Saeed A, Aslam MA, Younas F, Nawaz M, Mehnaz S and Fatima T, 2023. Diagnostic tools for zoonotic infections. In: Abbas RZ, Hassan MF, Khan A and Mohsin M (eds), Zoonosis, Unique Scientific Publishers, Faisalabad, Pakistan, Vol. 2: 432-445. <https://doi.org/10.47278/book.zoon/2023.193>

CHAPTER HISTORY

Received: 25-Jan-2023 Revised: 27-May-2023 Accepted: 21-June-2023

^{1,3,4, 5,7,8}Department of Chemistry, Government College University Faisalabad, Faisalabad

²Department of Biochemistry, Government College University Faisalabad, Faisalabad

⁶Department of Applied Chemistry, Government College University Faisalabad, Faisalabad

⁹Department of Parasitology, Faculty of Veterinary Science, University of Agriculture, Faisalabad

¹⁰Department of Pathobiology, Riphah College of veterinary sciences (RCVetS), Lahore Campus, Riphah International University

*Corresponding author: toobabatoolpakist@gmail.com


1. INTRODUCTION

Some unidentified and mysterious illnesses with no proper treatment have been observed all over the globe (He et al. 2020; Smolarz et al. 2021; Shafaati and Zandi 2022). Table 1 shows rapid diagnosis for molecular and non-molecular zoonotic infections.

Furthermore, such diseases are unique to specific regions of the world and have spread to new ones (Malki et al. 2020). Therefore, we need to be able to spot emerging diseases and determine its cause (Connolly et al. 2021).

Table 1: Rapid diagnosis for molecular and non-molecular zoonotic infections.

Molecular diagnostic methods	Non-molecular diagnostic methods
<ul style="list-style-type: none"> • Polymerase chain reaction (PCR) • Hybridization • Metagenomics • Gene probing • DNA Fingerprinting • rRNA sequencing • DNA Base composition • Electron Microscope • Mass Spectrometry • Restriction Endonuclease and Oligonucleotide Fingerprinting 	<ul style="list-style-type: none"> • Cultures • Beta-Glucan Assay • CAGTA Assay • Histopathology • Glutomanan assay • Serological Tests (Antigen tests, Antibody tests) • Optical microscope



2. DIVERSE APPROACHES

The technique that incorporates simultaneous and diverse approaches is the most promising for evaluating novel diseases (Basith et al. 2020).

2.1. VIRAL METAGENOMICS

A culture-independent method called viral metagenomics is used to examine all viral genetic populations in biological material (Sommers et al. 2021). This methodology becomes a powerful tool for identifying

ZOONOSIS

new and emerging viruses, considering that animals remain a reservoir for the virus that can cause zoonosis. Less than 1% of bacterial hosts have been cultivated, so it is difficult to identify and measure the dynamics of the viral population in the environment (Zhang et al. 2022). On the whole, these restrictions can be overcome by metagenomics investigation of uncultured viral communities, which can also offer insights into the makeup and organization of environmental viral communities.

2.2. CLINICAL RECOGNITION

An immediate individual or group recognizes a disorder that does not share traits with any other recognized condition (Shah et al. 2020). It could be an illness with a recently identified or previously documented ailment with an atypical presentation. As a result, it is advantageous to alert hospital infectious disease specialists, emergency room physicians, and other combatants to the potential for a novel disease or one that has not previously been observed in the region (Sami et al. 2021).

2.3. STANDARD LABORATORY ASSESSMENTS

It could be an illness with a recently identified or previously documented ailment with an atypical presentation. Therefore, it is advantageous to alert hospitals for novel disease specialists, emergency room physicians, and other combatants to the potential for a novel disease or one not previously observed in the region to encourage such recognition.

2.4. EPIDEMIOLOGY STUDIES

The epidemiological field is a scientific, systematic, and analytical study of the frequency, distribution, and origins of medical symptoms and events in particular groups (such as communities, schools, cities, regions, nations, and the world) (Luca et al. 2022).

The investigations may shed light on the new condition's distribution, the race, age, and sex of vulnerable people, the immune response of those who are resilient and those who are potential animal vectors or recipients, the incubating period, and the mode of transmission (Belbasis and Bellou 2018).

2.5. LABORATORY TESTS

All conceivable laboratory tests need to be performed when an indication of a new and significant disease is indicated. These exams comprise of:

- Microscopy
- Electron microscopy
- Culture
- Measurement of antibodies serum
- Antigen capture assays
- Nucleic acid amplifications
- Mass spectrometer
- Genes Measurements
- Vaccination of test subjects

2.6. MICROSCOPY

A microscope can be used for a variety of things, depending on its type. While some are utilized in instructional settings, others are appropriate for biological purposes (DeVree et al. 2021).

ZOONOSIS

Light microscopy can be performed quickly, but precision depends on the equipment's quality and the microscopist's competence. The ability of doctors to employ microscopy for diagnosis outside of an accredited laboratory is frequently constrained by regulations. To differentiate between invasive illness and surface colonization—a distinction that is difficult to make using culture methods—microscopic inspection of tissue may be necessary (Richert-Poggeler et al. 2019).

3. MICROSCOPY USING ELECTRONS AND IMMUNOELECTRON MICROSCOPY

These techniques are best suited for quickly detecting viral isolates from cell cultures and actual specimens (Madanayake et al. 2023). Electron microscopy only allows for family-level identification, whereas immunoelectron microscopy with a suitable, particular antibody may allow for the creation of finer differences (Gulati et al. 2019).

4. CULTURE

Culture is the development of microorganisms located in a nutrient-rich either liquid or solid medium; more organisms make identification easier. Antimicrobial susceptibility testing is also made more accessible by culture (Namdari et al. 2019).

5. IMMUNOLOGIC TESTS

The antigens are used to search for pathogen-specific antibodies in the patient's samples. Detection antibody for a pathogen's antigen in the patient's samples. Although there are different protocols for handling specimens, they should normally be chilled or frozen if testing needs to be postponed to avoid bacterial contamination proliferation (Normann et al. 2020). There are numerous techniques used or put forth for the cultivation-independent characterization of infectious pathogens (Gilboa et al. 2020; Preena et al. 2020). Through the use of molecular pathogen discovery techniques, new pathogens connected to both acute and chronic illnesses have been discovered during the past ten years (Belizario et al. 2021).

6. LIMITATIONS

6.1. THE PATHOGEN'S HOST RANGE

A pathogen once exclusive to dogs, swine, or cats now threatens human life (Tomori and Oluwayelu 2023).

6.2. MODIFICATION OF TISSUE TROPISM

A virus that in the past only occasionally caused moderate enteritis now frequently causes severe myocarditis and severe encephalitis (Shieh 2022).

6.3. IMMUNE EVASION

Large DNA viruses with many genes that help them escape or manipulate the host's immunity include orthopox viruses (Verdonck et al. 2022). These genes could undergo mutations that alter the virus's host range. Mutations can cause viruses of all sizes, including RNA viruses, to switch species. Additionally, some

ZOONOSIS

RNA fragments are capable of recombining to create new variations that are immune-evading for both humans and the animals that serve as their reservoirs (Ma et al. 2019).

6.4. ENVIRONMENTAL EFFECTS

Human beings' activities include constructing routes across the bush and moving animals or vectors (Mishra et al. 2021).

6.5. ZOONOSES WITHOUT A VECTOR

Pathologists can now identify the root cause of dying more quickly than ever before and frequently when it would have been impossible in the past, using techniques like polymerase chain reaction (PCR) testing, in situ hybridization, and immunohistochemistry (Sledzinska et al. 2021).

Table 2: Main categories of tests used to determine an accurate diagnosis of a viral outbreak in an animal

Taxonomy	Techniques	Examples	References
Family (genus)	<ul style="list-style-type: none"> Complement Fixation Electron Microscopy Cytopathology 	GBV-A, a virus from the family Flaviviridae, is found in wild monkeys, whereas GBV-B, from the genus Hepacivirus, is found in infected tamarins. Pegivirus is the name of the fourth genus in the Flaviviridae GBV-D and GBV-C family observed in bats, rodents, horses, and pigs	(Koonin et al. 2021)
Species (types)	Neutralization	Brown bats' <i>Eptesicus fuscus</i> and <i>Myotis lucifugus</i> were examined for the presence of the rabies virus and antibodies that can neutralize it.	(Gold et al. 2020)
Subtype	Kinetic neutralization Monoclonal antibody serology	Tembusu virus (TMUV), a new flavivirus that is developing in ducks, was neutralized by an antibody response in 2010. Influenza-neutralizing antibodies A virus and the effectiveness of their preventative measures in mice	(Lv et al. 2019)
Variant	<ul style="list-style-type: none"> Restriction Endonuclease Oligonucleotide fingerprint Nucleic acid hybridization 	Dromedaries with MERS-CoV Infection variant detection are detected and in pregnant white-tailed deer, SARS-CoV-2 and its alpha form were found and also in Norway rats. Endonuclease fragments were used to find the <i>bla</i> CTX-M-15 and <i>bla</i> CTX-M-1 (CTX-M ESBL gene variants) in <i>Escherichia coli</i> isolates and animal faeces	(Lado et al. 2021; Cool et al. 2022; Higgins et al. 2023)
Mutant (including by point mutation)	Nucleic acid sequencing	Asiatic lions in India were infected with the SARS-CoV-II Delta mutation	(Karikalan et al. 2022)

7. VIRAL DIAGNOSIS

There are three main categories of tests used to determine an accurate diagnosis of a viral outbreak in an animal:

1. Characterization and isolation of the causative virus.
 2. Measurement and detection of antibodies.
 3. Direct demonstration of viral nucleic acids, viral antigens, virions, or in tissues, secretions and excretions.
- There are numerous interesting diagnostic tests employing nanomaterials that are focused on the chemicals that cause animal diseases, but many of them have not yet reached a balance between specificity, sensitivity, cost-effectiveness, and repeatability. (Ramakrishnan et al. 2021). Table 2 shows main categories of tests used to determine an accurate diagnosis of a viral outbreak.

7.1. SPECIMEN PREPARATION FOR INOCULATION

Once the material has arrived at the laboratory, it should be handled and immunized as soon as feasible. If delays of more than a day are anticipated, the specimen may be frozen. Swabs are treated by rotating them in the vessel of transport and applying significant pressure to the container's side to release the substance (MacDonald et al. 2022). The excrement is dispersed using a vortex mixer. Tissue samples are meticulously cut with scissors and homogenized in a glass or mechanical homogenizer.

Table 3: The main serological methods in diagnostic virology.

Techniques of choice	Working Principle	Examples	References
Virus Neutralization	Antibody prevents cytopathology, protects animals, or decreases by neutralizing the virion's infectivity.	Antibodies against the West Nile virus were discovered in wild birds from Southern Spain. Virus-neutralizing antibodies and related variables against rabies in the immunized domestic dogs of Kathmandu Valley, Nepal.	(Ferraguti et al. 2016) (Rimal et al. 2020)
Hemagglutination Inhibition	Antibody suppresses hemagglutination of viruses.	the In Kerman, Iran's southeast, dogs were used in an experiment to test for the avian H ₉ N ₂ influenza virus's ability to hemagglutinate.	(Saberi et al. 2019)
Complement Fixation	Complement is bound by the antigen-antibody complex, rendering it effective to lyse hemolysis-sensitive sheep red blood cells.	Complement fixation test for detection of specific antibodies against bovine brucellosis in chosen peasant association in Guto Gida district, East Wollega Zone, Oromia Ethiopia	(Tosisa et al. 2020).
Immunoelectron Microscopy	Electron microscopy can reveal virions that have aggregated into antibodies.	After a single dose of immunization, immunoelectron microscopy-based immunogenicity of a recombinant VSV-vectored SARS-CoV vaccine generated strong protection in rhesus monkeys.	(Shan et al. 2022)
Immunofluorescence	The fluorescent antibody binds to subcellular antigens and fluoresces under ultraviolet light microscopy.	To investigate the seroprevalence of <i>Bartonella henselae</i> in Egyptian cats and people, an immunofluorescent test was performed	(Sayed et al. 2022)
Immunoperoxidase Staining	The intracellular antigen is recognized by the peroxidase-labeled antibody, and upon the addition of substrate, colored precipitate results.	<i>Brucella melitensis</i> in experimentally infected fetuses is diagnosed by immuno-peroxidase staining in the liver of the foetus.	(Mazlan et al. 2021)
ELISA (Enzyme-linked immunosorbent assay)	The reaction causes the substrate to change colour when an antibody or antigen that has been enzyme-labeled binds to it.	To find antibodies against <i>Toxoplasma gondii</i> in a cohort of hunting dogs, researchers used an ELISA with the modified agglutination test (MAT)	(Bellatreche et al. 2022)
Radioimmunoassay	An antigen or antibody that has been radiolabeled binds to it, such as when it is connected to the solid phase.	The ELISA test was developed based on radioimmunoassay to test samples from potentially infected animals.	(Broussard 2020)
Immunodiffusion	In a gel, soluble antigens and antibodies precipitate in clear lines.	Agar gel immunodiffusion assay for the diagnosis of canine brucellosis using <i>Brucella ovis</i> antigen	(Bolotin et al. 2022)

ZOONOSIS

Membrane filters with an average pore diameter of 0.45µm filter out contaminating microorganisms before inoculation. Then mycoplasmas can pass through these kinds of filters. The suspension needs to be refiltered using 0.22µm filters to get rid of mycoplasmas after the virus has been successfully isolated and grown to a significant titer. Faeces and tissue homogenates should be diluted by at least a factor of 10 before being centrifuged at 1000g for 15 minutes to produce a filterable supernate. The virus concentration is believed to be very low, so a high dosage of antibiotics may be chosen for filtration (Diaz-Linan et al. 2021).

7.2. UTILIZING SEROLOGY TO IDENTIFY VIRAL ISOLATES

The main serological methods used to identify various viruses have been enlisted in Table 3.

7.3. COMPLEMENT FIXATION

Indirect diagnostic tests with great sensitivity like the complement fixation test (CFT) have a lot of false-positive outcomes (Elschner et al. 2021).

Inactivated serum samples are diluted twice in Complement fixation test (CFT) buffer and combined with CFT antigen and complement hemolytic units. Sera, complement, and antigen are combined in the plates and left to interact overnight at 4°C. This allows the complement to be fixed.

The plates are then incubated for 45 minutes at 37°C with a 2% suspension, then centrifuged for 5 minutes at 600g, and then the process is repeated. Samples exhibiting 25-75% hemolysis are regarded as suspicious, samples showing 100% hemolysis in a dilution are classified as negative. The classification of all suspicious test results is affirmative (Elschner et al. 2019).

Complement fixation can be utilized to classify an isolate and perform initial screening on it. Essentially a slightly streamlined version of the complement fixation of complement fixation test, the immune-adherence hemagglutination test is currently employed more commonly to identify antibodies than antigens (Coombs 2012).

7.4. HEMAGGLUTINATION AND INHIBITION OF HEMAGGLUTINATION

Numerous viral families' virions attach to red blood cells to hemagglutinate them. Hemagglutination is suppressed if antibodies and viruses are combined before the inclusion of red blood cells (Cho et al. 2022). Dissociating the virions using detergents can boost the hemagglutination level of some viruses, such as the canine distemper virus (Suarez et al. 2020). Antisera may need to be prepared to get rid of hemagglutination inhibitors that aren't specific. Test for determining the titer of antibodies to viral hemagglutinin by inhibiting hemagglutination. In titers, dilutions are stated as reciprocals (Spackman and Sitaras 2020).

8. METHODOLOGY

1. Periodate is applied to the sera, which is subsequently subjected to heat at 56°C for a half-hour to inactivate non-specific hemagglutination inhibitors.
2. After that, the sera are diluted in microtiter wells. The appropriate viral strain is then injected with hemagglutinating units into each well. Each well receives 0.05 ml of red blood cells following a 30-minute room temperature incubation period (Chivukula et al. 2021; Zhang et al. 2022).
3. Hemagglutination has been prevented where there is enough antibody to coat the virions, and as a result, at the bottom of the well, the erythrocytes condense to form a button (Saegerman et al. 2021).

ZOONOSIS

Erythrocytes get agglutinated by the virus when there is insufficient antibody present. For example, the horse lacked any antibodies that could have blocked the hemagglutinin-inducing effects of this particular influenza virus strain. The first dose of the vaccine resulted in some antibody production, and the second dose induced a booster response (Drozd et al. 2020).

8.1. NEUTRALIZATION OF VIRUS

A unique antibody may neutralize a virus's infectiousness through several different processes. To get rid of general viral inhibitors. The serum first needs to be inactivated by being heated at 56°C for 30 minutes (Cuevas et al. 2022). Suitable cell cultures are infected with serum-virus combinations. Afterward, they are allowed to develop until the virus-only controls show cytopathic effects by reducing a virus's ability to infect, an antibody shields cells against viral annihilation (Amanat et al. 2020).

8.2. RESTRICTION ENDONUCLEASE AND OLIGONUCLEOTIDE FINGERPRINTING

For the majority of routine diagnostic purposes, even to the level just mentioned, it is often not necessary to isolate antigenically. However, there are circumstances in which going further to identify differences between variants or subtypes within a particular serotype could provide viral epidemiological data. Similarly, viral DNA can be isolated from infected cells or virions, and the fragments can then be separated using agarose gel electrophoresis after being cut with properly chosen restriction endonucleases (Guo et al. 2022).

Ethidium bromide per silver staining is required to obtain restriction endonuclease fragment patterns, which are frequently referred to as fingerprints. All dsDNA virus families have found a use for methodology, particularly referred to as fingerprints. All dsDNA virus families have found use for the methodology, particularly in epidemiology research but also in pathogenesis research. These methods' resolution allows us to distinguish different isolates of the same viral species, even if they did not all come from the same epizootic, depending on the viral family. Minor amounts of genetic drift, which are typically not reflected in serological differences, can occasionally be detected using this technique (Laudermilch and Chandran 2021).

8.3. ELECTRON MICROSCOPY FOR THE DIRECT DETECTION OF VIRIONS

Electron microscopes are used for quick viral diagnosis. Negative staining techniques and understanding the concentration of virions are crucial for accuracy. Technology in the medical field is constantly advancing. This method can diagnose viral skin disease using vesicular fluid, scrapings, or scabs. It can also be used for accurate diagnosis in cell culture (Hopfer et al. 2021).

8.4. IMMUNOELECTRON MICROSCOPY

It is a technique that uses immune serum to increase the susceptibility of electron microscopic techniques. After mixing the sample with the antibody for an overnight period, the sample is typically cleared by low-speed centrifugation.

The immune complexes are subsequently centrifuged at a low speed to form pellets, which are then negatively stained. The antibody used could be a mixture of these antibodies, serum from an old animal that is hyperimmune to many viruses, or type-specific monoclonal or polyclonal antibodies (Zhang et al. 2022).

8.5. DIRECT IDENTIFICATION OF VIRAL ANTIGENS

These techniques rely on the direct interaction of viral particles, or antigens, with specific antibodies that have been pre-labeled in some way to quickly identify the interaction, in situ in tissues, excretions, or secretions. The labeling techniques used—are immunoperoxidase staining and radioimmunoassay.

The labeling methods are divided into four categories: radioimmunoassay, immunoperoxidase staining, immunofluorescence, and ELISA. Viral antigens can also be identified using two proven serological techniques, precipitation and complement fixation (Bassani-Sternberg et al. 2016).

8.6. IMMUNOFLUORESCENCE

Immunofluorescence is a method of unique significance in the quick identification of viral infections due to its specificity, speed, relative simplicity, and sensitivity (Zhang et al. 2022).

The classic instance of immunofluorescence is the diagnosis of rabies, for which immunofluorescence has been recognized as diagnostic for more than 20 years (Chiebao et al. 2019).

9. STAINING WITH IMMUNOPEROXIDASE

An alternate approach involves using an enzyme-labeled antibody to locate and detect viral antigens in infected cells. The process results in a pore-durable, non-fading, and anatomically clearer preparation than immunofluorescence, and it uses less expensive equipment. Similar steps and ideas apply to immunofluorescence (Burrell et al. 2017).

The conjugated antibody, bound to the antigen through a direct or indirect procedure, is detected by adding a substrate suited for the specific enzyme. In the case of peroxidase, this is H_2O_2 coupled with a benzidine derivative. Endogenous peroxidase, which is found in many tissue cells, especially leukocytes, causes false positive results, which is a drawback of the method. By using a diligent approach and proper controls, this issue can be avoided (Arshad et al. 2022).

9.1. RADIOIMMUNOASSAY

A radioactive element, most frequently ^{125}I , serves as the label in radioimmunoassay. The technique is incredibly sensitive, allowing the detection of viral antigens at low concentrations ranging from 10–12M. There are numerous radioimmunoassay techniques available. The principles are the same for both direct and indirect approaches as for immunofluorescent staining.

In the most basic configuration, a sample that might contain a virus or viral antigen is allowed to attach to the bound antibody, washed, and then an antiviral antibody is measured in a gamma counter after additional washing (Burrell et al. 2017).

A more popular approach is indirect radio-immunoassays, which include a second layer of ^{125}I -labelled anti-IgG as an indicator antibody in place of the detection antibody. Different animal species must be used to generate the antiviral antibodies that make up the capture and detection antibodies (Inoue et al. 2010).

9.2. DIRECT ISOLATION OF LEPTOSPIRES FROM CLINICAL SAMPLES

A common method for identifying *Leptospira* directly from clinical samples is dark-field microscopy (DFM). Early detection of leptospires in bacterial infections is crucial. The direct fluorescent antibody (DFM) is

ZOONOSIS

highly sensitive for detecting leptospire in both blood and CSF, with a sensitivity rate of 64.7%. DFM is a good way to diagnose the early stages of an illness, with high sensitivity in identifying leptospire in blood and CSF. A skilled specialist is needed and 100 fields on each slide must be studied to deem it negative. As the illness progresses, DFM's sensitivity may decrease.

9.3. LEPTOSPIRES' CULTURE

A Bunsen burner will be used to aseptically add blood samples dropwise to the semi-solid medium that has been prepared. Within 24 hours, the sample-containing medium will be switched over to the liquid media. Every day for three months, each tube will be examined for the development of leptospire, and after every 2-3 weeks, they will be routinely switched to a new medium. The media developed by Stuart, Korthoff, and Fletcher is also utilized for culture. Although cultivation is the most common method of detection, its application for routine isolation is debatable because it requires a longer period for development, with a gestation period of 6–20 hours (Hornsby et al. 2020).

9.4. SEROLOGICAL METHODS

The serum of people who have contracted *Leptospira* can be tested using a variety of serological assays for antibodies against the parasite. We can detect IgG and IgM antibodies with MAT and ELISA-based test kits, and a wide variety of commercial fast diagnostic card tests that are already on the market (Trozzi et al. 2023).

9.5. MOLECULAR TECHNIQUES

Molecular methods can quickly and accurately diagnose infections, including leptospire, in medical or environmental samples (Girones et al. 2010).

9.6. POLYMERASE CHAIN REACTION (PCR)

Due to its specificity, sensitivity, and capacity to identify even the smallest amounts of nucleic acid particles. PCR is the best way to identify infections. Quantitative PCR is better than MAT or culture for quick and reliable results (Wood et al. 2019). This finding has important implications for medical and epidemiological analyses of this worldwide neglected disease. *Leptospira* is also detected using a variety of PCR techniques, including nested PCR, randomly primed PCR, etc. The drawbacks of PCR are similar to those of other diagnostic techniques and include complex lab requirements, degradation, false-positive results, initial validation, etc. These problems necessitate highly competent and trained personnel as well as a significant financial commitment (Hoorfar et al. 2004).

9.7. QUANTITATIVE PCR

PCR techniques detect *Leptospira*, but have limitations such as specialized equipment, sample degradation, and false positives. qPCR (real-time PCR) is preferred for accurate results in a short time by monitoring the DNA amplification rate. To evaluate disease severity, bacterial DNA amount and density must be assessed, often through targeting the *LipL32* gene. qPCR is a reliable method for the early detection of leptospirosis but is challenging to implement in low-resource labs due to the required skilled personnel and expensive equipment (Ruijter et al. 2021).

9.8. IMMUNOCAPTURE-POLYMERASE CHAIN REACTION

New techniques have been developed to quickly identify leptospire from clinical samples, including the immunocapture-polymerase chain reaction methods created in 2018 using ELISA and PCR. A combination of molecular and serological methods is important for an accurate diagnosis of leptospire. The IC-PCR technique is a powerful tool that rapidly identifies leptospire and provides crucial information about specific serovars or serogroups present. This study shows it to be more accurate than traditional PCR methods (Jian et al. 2020).

9.9. OTHER MOLECULAR METHODS

Other molecular methods such as RAPD (random amplified polymorphic DNA), nucleic acid probe techniques, DNA-DNA hybridization, REA (restriction enzyme analysis), fingerprinting, PFGE (pulsed-field gel electrophoresis), etc., are available in highly developed laboratory settings. They support research into and comprehension of genetic diversity and genomic diversity profiles.

Several diagnostic techniques are available to detect and diagnose Leptospirosis, including ELA (IgM-Enzyme Immunoassay), MCAT (microcapsule agglutination test, LEPTO Dipstick, macroscopic SAT, IHA (indirect haemagglutination assay), and LEPTO Dri-dot. However, these rapid tests are less sensitive and specific than IgM rapid and immunochromatography techniques, despite the availability of several serological rapid tests.

CONCLUSION

Our ability to diagnose the cause of the disease is greatly improved by following all the stages consistently and sequentially. This method not only increases the scientific knowledge base but also helps zoo employees conduct retrospective studies and aids other researchers with the new database. It also enhances the ability of the zoological community to recognize disease trends and enables researchers to identify diseases that pose a threat to captive and wild animals.

REFERENCES

- Amanat et al., 2020. An in vitro microneutralization assay for SARS-CoV-2 serology and drug screening. *Current Protocols in Microbiology* 58(1): Article # e108.
- Arshad et al., 2022. Nanotechnology for therapy of zoonotic diseases: a comprehensive overview. *Chemistry Select* 7(21): Article # e202201271.
- Basith et al., 2020. Machine intelligence in peptide therapeutics: A next-generation tool for rapid disease screening. *Medicinal Research Reviews* 40(4): 1276-1314.
- Bassani-Sternberg et al., 2016. Direct identification of clinically relevant neoepitopes presented on native human melanoma tissue by mass spectrometry. *Nature Communications* 7(1): Article # 13404.
- Belbasis L and Bellou V, 2018. Introduction to epidemiological studies. *Genetic Epidemiology: Methods and Protocols* 4: 1-6.
- Belizario et al., 2021. Breath biopsy and discovery of exclusive volatile organic compounds for diagnosis of infectious diseases. *Frontiers in Cellular and Infection Microbiology* 10: Article # 564194.
- Bellatreche et al., 2022. Comparison of a Commercial Enzyme-Linked Immunosorbent Assay (ELISA) with the Modified Agglutination Test (MAT) for the Detection of Antibodies against *Toxoplasma gondii* in a Cohort of Hunting Dogs. *Animals* 12(20): Article # 2813.
- Bolotin et al., 2022. Use of *Brucella Ovis* Antigen for Canine Brucellosis Diagnostics by Agar Gel Immunodiffusion Assay. In *Brucellosis 2022 International Research Conference* 78: 6-02.

- Broussard E, 2020. Helping to tackle the spread of zoonotic disease: The VETLAB Network. *IAEA Bulletin* 11: 56-59.
- Burrell et al., 2017. Laboratory diagnosis of virus diseases. *Fenner and White's Medical Virology* 135: 33-39.
- Chiebao et al., 2019. Congenital transmission of *Toxoplasma gondii* after experimental reinfection with Brazilian typical strains in chronically infected sheep. *Frontiers in Veterinary Science* 6: Article # 93.
- Chivukula et al., 2021. Development of multivalent mRNA vaccine candidates for seasonal or pandemic influenza. *NPJ Vaccines* 6(1): Article # 153.
- Cho et al., 2022. Broccoli Leaves Attenuate Influenza A Virus Infection by Interfering with Hemagglutinin and Inhibiting Viral Attachment. *Frontiers in Pharmacology* 13: Article # 2149.
- Connolly et al., 2021. Extended urbanization and the spatialities of infectious disease: Demographic change, infrastructure, and governance. *Urban Studies* 58(2): 245-263.
- Cool et al., 2022. Infection and transmission of ancestral SARS-CoV-2 and its alpha variant in pregnant white-tailed deer. *Emerging Microbes & Infections* 11(1): Article # 95-112.
- Coombs R, 2012. Assays Utilizing Red Cells. *Immunoassays for the 80s* 17: Article # 11.
- Cuevas et al., 2022. An In Vitro Microneutralization Assay for Influenza Virus Serology. *Current Protocols* 2(7): Article # e465.
- DeVree et al., 2021. Current and future advances in fluorescence-based visualization of plant cell wall components and cell wall biosynthetic machinery. *Biotechnology for Biofuels* 14(1): Article # 78.
- Diaz-Linan et al., 2021. Unmodified cellulose filter paper, a sustainable and affordable sorbent for the isolation of biogenic amines from beer samples. *Journal of Chromatography A* 1651: Article # 462297.
- Drozd et al., 2020. Obligate and facultative anaerobic bacteria in targeted cancer therapy: Current strategies and clinical applications. *Life Sciences* 261: Article # 118296.
- Elschner et al., 2019. Evaluation of the comparative accuracy of the complement fixation test, Western blot, and five enzyme-linked immunosorbent assays for serodiagnosis of glanders. *PLoS One* 14(4): Article # e0214963.
- Elschner et al., 2021. Validation of a Commercial Glanders ELISA as an Alternative to the CFT in International Trade of Equidae. *Frontiers in Veterinary Science* 8: Article # 628389.
- Ferraguti. 2016. West Nile virus-neutralizing antibodies in wild birds from southern Spain. *Epidemiology & Infection* 144(9): 1907-1911.
- Gilboa et al., 2020. Single-molecule analysis of nucleic acid biomarkers—A review. *Analytica Chimica Acta* 1115: 61-85.
- Girones et al., 2010. Molecular detection of pathogens in water—the pros and cons of molecular techniques. *Water Research* 44(15): 4325-4339.
- Gold et al., 2020. Rabies virus-neutralizing antibodies in healthy, unvaccinated individuals: What do they mean for rabies epidemiology. *Neglected Tropical Diseases* 14(2): Article # e0007933.
- Gulati et al., 2019. Immunoelectron microscopy of viral antigens. *Current Protocols in Microbiology* 53(1): Article # e86.
- Guo et al., 2022. Restriction-Assembly: A Solution to Construct Novel Adenovirus Vector. *Viruses* 14(3): Article # 546.
- He et al., 2020. Coronavirus disease 2019: What we know. *Journal of medical virology* 92(7): 719-725.
- Higgins et al., 2023. Portable Differential Detection of CTX-M ESBL Gene Variants, bla CTX-M-1, and bla CTX-M-15, from *Escherichia coli* Isolates and Animal Fecal Samples Using Loop-Primer Endonuclease Cleavage Loop-Mediated Isothermal Amplification. *Microbiology Spectrum* 11(1): Article # e03316.
- Hoorfar et al., 2004. Practical considerations in the design of internal amplification controls for diagnostic PCR assays. *Journal of Clinical Microbiology* 42(5): 1863-1868.
- Hopfer et al., 2021. Hunting coronavirus by transmission electron microscopy—a guide to SARS-CoV-2-associated ultrastructural pathology in COVID-19 tissues. *Histopathology* 78(3): 358-370.
- Hornsby et al., 2020. Isolation and propagation of leptospire at 37 °C directly from the mammalian host. *Scientific Reports* 10(1): 1-11.
- Inoue et al., 2010. Evaluation of a dengue IgG indirect enzyme-linked immunosorbent assay and a Japanese encephalitis IgG indirect enzyme-linked immunosorbent assay for diagnosis of secondary dengue virus infection. *Vector-Borne and Zoonotic Diseases* 10(2): 143-150.
- Janke et al., 2001. Paramyxovirus infection in pigs with interstitial pneumonia and encephalitis in the United States. *Journal of Veterinary Diagnostic Investigation* 13(5): 428-433.

- Jian et al., 2020. Quantitative PCR provides a simple and accessible method for quantitative microbiota profiling. *PloS One* 15(1): Article # e0227285.
- Karikalan et al., 2022. Natural infection of Delta mutant of SARS-CoV-2 in Asiatic lions of India. *Transboundary and emerging diseases* 69(5): 3047-3055.
- Koonin et al., 2021. The healthy human virome: From virus–host symbiosis to disease. *Current Opinion in Virology* 47: 86-94.
- Lado et al., 2021. Innate and adaptive immune genes associated with MERS-CoV infection in dromedaries. *Cells* 10(6): Article # 1291.
- Laudermilch E and Chandran K, 2021. MAVERICC: marker-free vaccinia virus engineering of recombinants through in vitro CRISPR/Cas9 Cleavage. *Journal of Molecular Biology* 433(9): 166896.
- Luca et al., 2022. Inequality, crime, and public health: A survey of emerging trends in urban data science. *Arxiv Preprint Arxiv 2212*: Article # 07676.
- Lv et al., 2019. Detection of neutralizing antibodies to Tembusu virus: implications for infection and immunity. *Frontiers in Veterinary Science* 6: Article # 442.
- Ma et al., 2019. In vitro and in vivo efficacy of a Rift Valley fever virus vaccine based on pseudovirus. *Human Vaccines & Immunotherapeutics*.
- MacDonald et al., 2022. *Pagana's Canadian Manual of Diagnostic and Laboratory Tests-E-Book*. Elsevier Health Sciences.
- Madanayake et al., 2023. Electron microscopic methods for virus diagnosis. In: Dhara AK, Nayak AK, editors. *Viral Infections and Antiviral Therapies*: Academic Press; pp: 121-140.
- Malki et al., 2020. Association between weather data and COVID-19 pandemic predicting mortality rate: Machine learning approaches. *Chaos, Solitons & Fractals* 138: Article # 110137.
- Mazlan et al., 2021. Pathological changes, distribution, and detection of *Brucella melitensis* in foetuses of experimentally-infected does. *Veterinary Quarterly* 41(1): 36-49.
- Mishra et al., 2021. Linkages between environmental issues and zoonotic diseases: with reference to COVID-19 pandemic. *Environmental Sustainability* 4(3): 455-467.
- Namdari et al., 2019. Comparative study of cultures and next-generation sequencing in the diagnosis of shoulder prosthetic joint infections. *Journal of Shoulder and Elbow Surgery* 28(1): 1-8.
- Normann et al., 2020. Pathogen-specific antibody profiles in patients with severe systemic infections. *Eur Cells & Mater* 39: 171-182.
- Preena et al., 2020. Unravelling the menace: detection of antimicrobial resistance in aquaculture. *Letters in Applied Microbiology* 71(1): 26-38.
- Ramakrishnan et al., 2021. Nanotechnology-based solutions to combat zoonotic viruses with special attention to SARS, MERS, and COVID-19: Detection, protection and medication. *Microbial Pathogenesis* 159: Article # 105133.
- Richert-Poggeler et al., 2019. Electron microscopy methods for virus diagnosis and high-resolution analysis of viruses. *Frontiers in Microbiology* 9: Article # 3255.
- Rimal et al., 2020. Detection of virus-neutralizing antibodies and associated factors against rabies in the vaccinated household dogs of Kathmandu Valley, Nepal. *PloS One* 15(4): Article # e0231967.
- Ruijter et al., 2021. Efficiency correction is required for accurate quantitative PCR analysis and reporting. *Clinical Chemistry* 67(6): 829-842.
- Saberi et al., 2019. Serological prevalence of avian H9N2 influenza virus in dogs by hemagglutination inhibition assay in Kerman, southeast of Iran. In *Veterinary Research Forum* 10(3): 249-250.
- Saegerman et al., 2021. Formal estimation of the seropositivity cut-off of the hemagglutination inhibition assay in field diagnosis of influenza D virus in cattle and estimation of the associated true prevalence in Morocco. *Transboundary and Emerging Diseases* 68(3): 1392-1399.
- Sami et al., 2021. A comprehensive review on global contributions and recognition of pharmacy professionals amidst COVID-19 pandemic: moving from present to future. *Future Journal of Pharmaceutical Sciences* 7(1): 1-16.
- Sayed et al., 2022. Serological and Molecular Detection of *Bartonella henselae* in Cats and Humans from Egypt: Current Status and Zoonotic Implications. *Frontiers in Veterinary Science* 9: Article # 859104.

- Shafaati M and Zandi M, 2022. Monkeypox virus neurological manifestations in comparison to other orthopoxviruses. *Travel Medicine and Infectious Disease* 49: Article # 102414.
- Shah et al., 2020. International Working Group on Transdiagnostic Clinical Staging in Youth Mental Health. Transdiagnostic clinical staging in youth mental health: a first international consensus statement. *World Psychiatry* 19(2): 233-242.
- Shan et al., 2022. Immunogenicity of a recombinant VSV-Vectored SARS-CoV vaccine induced robust immunity in rhesus monkeys after single-dose immunization. *Virologica Sinica* 37(2): 248-255.
- Sledzinska et al., 2022. Glioma 2021 WHO Classification: The Superiority of NGS over IHC in Routine Diagnostics. *Molecular Diagnosis & Therapy* 26(6): 699-713.
- Smolarz et al., 2021. Endometriosis: epidemiology, classification, pathogenesis, treatment and genetics (review of literature). *International Journal of Molecular Sciences* 22(19): Article # 10554.
- Sommers et al., 2021. Integrating viral metagenomics into an ecological framework. *Annual Review of Virology* 8: 133-158.
- Spackman E and Sitaras I, 2020. Hemagglutination inhibition assay. *Animal Influenza Virus: Methods and Protocols* 2020: 11-28.
- Suarez et al., 2020. Newcastle disease, other avian paramyxoviruses, and avian metapneumovirus infections. *Diseases of Poultry* 78: 109-166.
- Tomori O and Oluwayelu D O, 2023. Domestic animals as potential reservoirs of zoonotic viral diseases. *Annual Review of Animal Biosciences* 11: 33-55.
- Tosisa et al., 2020. Prevalence and antimicrobial susceptibility of *Salmonella* and *Shigella* species isolated from diarrheic children in Ambo town. *BMC Pediatrics* 20(1): 1-8.
- Torres-Sangiao et al., 2021. Application and perspectives of MALDI–TOF mass spectrometry in clinical microbiology laboratories. *Microorganisms* 9(7): Article # 1539.
- Trozzi et al., 2023. Comparison of Serological Methods for Tick-Borne Encephalitis Virus-Specific Antibody Detection in Wild Boar and Sheep: Impact of the Screening Approach on the Estimated Seroprevalence. *Viruses* 15(2): Article # 459.
- Verdonck et al., 2022. Viral manipulation of host cell necroptosis and pyroptosis. *Trends in Microbiology* 30(6): 593-605.
- Virtanen et al., 2021. Kinetics of neutralizing antibodies of COVID-19 patients tested using clinical D614G, B. 1.1. 7, and B 1.351 isolates in microneutralization assays. *Viruses* 13(6): Article # 996.
- Wood et al., 2019. A comparison of droplet digital polymerase chain reaction (PCR), quantitative PCR and metabarcoding for species-specific detection in environmental DNA. *Molecular Ecology Resources* 19(6): 1407-1419.
- Zhang et al., 2022. Hemagglutinin stalk-binding antibodies enhance the effectiveness of neuraminidase inhibitors against influenza via Fc-dependent effector functions. *Cell Reports Medicine* 3(8): 100718.
- Zhang et al., 2022. Beyond neutralization: Fc-dependent antibody effector functions in SARS-CoV-2 infection. *Nature Reviews Immunology* 34: 1-16.
- Zhang et al., 2022. Recent insights into aquatic viruses: Emerging and reemerging pathogens, molecular features, biological effects, and novel investigative approaches. *Water Biology and Security* 25: Article # 100062.
- Zhang et al., 2022. Advanced point-of-care testing technologies for human acute respiratory virus detection. *Advanced Materials* 34(1): Article # 2103646.